



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/48, 31/22	A1	(11) International Publication Number: WO 98/51277 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/US98/09729 (22) International Filing Date: 13 May 1998 (13.05.98) (30) Priority Data: 60/046,343 13 May 1997 (13.05.97) US 60/080,695 3 April 1998 (03.04.98) US 09/076,374 12 May 1998 (12.05.98) US (71) Applicant: CASE WESTERN RESERVE UNIVERSITY [US/US]; 10900 Euclid Avenue, Cleveland, OH 44106 (US). (72) Inventors: BRUNENGRABER, Henri; 16700 South Woodland, Shaker Heights, OH 44120 (US). BOMONT, Catherine; 193 Spruce Mill Lane, Scotch Plains, NJ 07076 (US). DAVID, France; 19826 Van Aken Boulevard, Shaker Heights, OH 44122 (US). HALLOWELL, Peter, T.; 1097 Pennfield, Cleveland Heights, OH 44121 (US). COOPER, Kevin, D.; 55 Quail Hollow Drive, Moreland Hills, OH 44022 (US). KASOUMOV, Takhar; 1898 East 123rd Street #3, Cleveland, OH 44106 (US). (74) Agent: MINNICH, Richard, J.; Fay, Sharpe, Beall, Fagan, Minnich & McKee, 7th floor, 1100 Superior Avenue, Cleveland, OH 44114-2518 (US).		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, GH, GM, GW, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: MEDICAL USES OF PYRUVATES (57) Abstract A pyruvate compound suitable for cosmetically or dermatologically administering to the skin and for use in treating diabetic ketosis or other medical treatments. The compound includes a pyruvate selected from the group of pyruvate thioester, dihydroxyacetone-pyruvate, and an ester of pyruvate and a sugar or a polyol.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MEDICAL USES OF PYRUVATES

Background of the Invention

This invention relates to several new pyruvate compounds and methods of (i) treating ischemia in mammalian hearts, lungs, veins, arteries and other organs or tissues, (ii) accelerating ethanol oxidation/preventing acute toxic effects of ethanol on the liver, (iii) using the novel pyruvates in other recognized uses of pyruvates including, but not limited to, treating of secondary effects of diabetes, lowering of cholesterol and lipid levels, as a nutrition source which can provide as much as 100% of caloric requirements and to treat injured organs requiring a readily accessible energy source, (iv) topical applications of pyruvates in the treatment of dermatological wounds or diseases and prevention thereof and to generally improve skin health, and (v) treatment of diabetic ketosis.

Description of the Art

Ischemia is defined herein as the interruption of oxygen supply, via the blood, to an organ or to part of an organ. Examples of ischemic events include (i) myocardial, cerebral, or intestinal infarction following obstruction of a branch of a coronary, cerebral, or mesenteric artery, and (ii) removal and storage of an organ prior to transplantation. In the case of myocardial infarction, prompt restoration of blood flow to the ischemic myocardium, i.e. coronary reperfusion, is a key component of the treatment. This is because mortality is directly related to infarct size (tissue necrosed) which is related to the severity and duration of the ischemic event.

Notwithstanding the need to supply an organ cut-off from a normal blood supply with oxygen, it has been found that reperfusion injury may occur upon restoration of blood flow. This results from the production of reactive oxygen species

-2-

(ROS), namely, hydrogen peroxide, hydroxyl radicals and superoxide radicals which are formed from both extracellular and intracellular sources. ROS are highly reactive species that, under normal conditions, are scavenged by endogenous defense mechanisms. However, under conditions of post-ischemic oxidative stress, ROS interact with a variety of cellular components, causing peroxidation of lipids, denaturation of proteins, and interstitial matrix damage, resulting in increase of membrane permeability and release of tissue enzymes.

In an attempt to minimize these undesirable side effects of perfusion, researchers Simpson, et al., (*Free Radical Scavengers and Myocardial Ischemia, Federation Proceedings, Volume 46, No. 7 May 15, 1987*) suggest the use of an inhibitor of ROS production to protect the reperfused myocardium. The Simpson, et al. disclosure is particularly directed to the use of agents and inhibitors (ex. allopurinol) that reduce ROS levels. In a similar context, Brunet, et al., (*Effects of Acetylcysteine, Free Radical Biology and Medicine, Volume XX, No. X 1995*) suggest the use of acetylcysteine to reperfuse hearts. In particular, the article concludes that acetylcysteine treatment decreases the production of ROS in reperfused rat hearts.

In a further effort directed to protecting reperfused heart tissue, United States Patent 5,075,210, herein incorporated by reference, discloses a process for reperfusing a heart for transplantation. The patent discloses a cardioplegic solution containing sodium chloride, potassium chloride, calcium chloride, sodium bicarbonate, sodium EDTA, magnesium chloride, sodium pyruvate and a protein.

United States Patent 5,294,641, herein incorporated by reference, is directed to the use of pyruvate to prevent the adverse effects of ischemia. The pyruvate is administered prior to a surgical procedure to increase a patient's cardiac output and heart stroke volume. The pyruvate is administered as a calcium or sodium salt. The pyruvate can alternatively be an ester of pyruvic acid such as ethylamino pyruvate.

-3-

Similarly, United States Patent 5,508,308, herein incorporated by reference, discloses the use of pyruvyl glycine to treat reperfusion injury following myocardial infarction.

5 United States Patent 4,988,515 and 5,705,210, herein incorporated by reference, use pyruvate salts in cardioplegic solutions and in preservation solutions for the heart before transplantation. United States Patent 4,970,143, herein incorporated by reference, discloses the use of acetoacetate
10 for preserving living tissue, including addition of the pyruvate to the preservation solution.

United States Patent 5,100,677 herein incorporated by reference, discloses the composition of various parenteral solutions. Of interest is a recommendation to include
15 pyruvate anions (apparently from metal salts) in intravenous solutions. In United States Patent 5,183,674, herein incorporated by reference, pyruvate compounds are used as foodstuff. United States Patent 5,134,162 herein incorporated by reference, focuses on the use of pyruvate to
20 lower cholesterol and lipid levels in animals. United States Patent 5,047,427, deals with the use of pyruvate for improving the condition of diabetics, while United States Patent 5,256,697 suggests the use of pyruvyl-aminoacid compounds, each of which is herein incorporated by reference.

25 In addition, United States Patent 5,283,260, herein incorporated by reference, is directed to treatment of diabetes with a physiologically acceptable form of pyruvate. The patent discloses a pyruvate compound in the form of a covalently linked pyruvyl-amino acid. By utilizing this type
30 of a pyruvate delivery system, the negative effect of pyruvate salt is avoided. However, administration of large amounts of pyruvate-amino acid may result in nitrogen overload which could harm patients with liver and/or kidney pathology.

35 Notwithstanding the acceptance of pyruvate as an effective component of a reperfusion solution or other varied applications, pyruvic acid is a strong and unstable acid which cannot be infused as such. Furthermore, it has been

-4-

recognized that traditional pharmacological pyruvate compounds, such as salts of pyruvic acid, are not particularly physiologically suitable. For example, these compounds lead to the accumulation of large concentrations of ions (ex. calcium or sodium) in the patient's body fluids. Similarly, amino acid compounds containing pyruvate can lead to excessive nitrogen loads. It has also been proposed to infuse pyruvylglycine, the amide function of which is presumably hydrolyzed in plasma and/or tissues, thus liberating pyruvate. However, at the high rates of pyruvylglycine infusion required to achieve 1 mM pyruvate in plasma, the glycine load may be harmful to patients suffering from hepatic or renal pathologies. Also, flooding plasma with glycine may interfere with the transport of some amino acids across the blood-brain barrier. Accordingly, while potentially suitable to organ preservation, these pyruvate compounds are less suited to treating an organ in vivo, and it is recognized that a need exists to provide a pyruvate delivery compound which is more physiologically acceptable.

In addition, pyruvates have been demonstrated to have several dermatological uses. For example, U.S. Patents 5,580,902; 5,602,183; and 5,614,561, herein incorporated by reference, disclose the use of pyruvate salts and ethyl ester to enhance therapeutic effects of cosmetics and pharmaceuticals. It is believed that the present pyruvate compounds are suited for at least all of the treatments identified in these patents and can be formulated into ointments, creams, etc., much in the same ways identified therein, and in other systems apparent to one skilled in the art.


Therefore, it is desirable in this field to have an alternate physiologically compatible therapeutic pyruvate compound. In this regard, the novel pyruvate compounds of this invention permit the use of pyruvate to treat ischemic events, ethanol poisoning, acetaminophen poisoning and other conditions recognized to be effectively treated with pyruvate because sufficiently high loads of pyruvate can be administered without a toxic constituent.

-5-


Summary of the Invention

One novel pyruvate compound of this invention comprises a pyruvate thioester. Preferably, the thiol is cysteine or homocysteine. In a particularly preferred form, the compound
 5 is a N-acetyl ethyl ester of the cysteine or homocysteine amino acid.

One preferred compound is:




A further novel compound of the present invention is a glycerol-pyruvate ester. A particularly preferred form of a
 10 glycerol-pyruvate ester will be of the formula:




and one or two R may be a short-chain acyl such as acetyl or propionyl.

20 and more preferably




30 Another novel compound of the present invention is a dihydroxyacetone-pyruvate ester. A particularly preferred form of this compound is of the formula:

-6-




and one R may be a short-chain acyl such as acetyl or propionyl.

10 and more preferably




20 Another novel compound of the present invention is a ribose pyruvate ester. A particularly preferred form of this compound is of the formula:



where R is H, pyruvyl, or a short-chain acyl such as acetyl or propionyl, not all four R's are H, one to four R's may be pyruvyl, and one to three R's may be short-chain acyl.

25 Another novel compound of the present invention is a glucose pyruvate ester. Particularly preferred forms of this compound (pyranose and furanose) are of the formulae:



30 where R is H, pyruvyl, or a short-chain acyl such as acetyl or propionyl, not all five R's are H, one to five R's may be pyruvyl, and one to four R's may be short-chain acyl.

-7-

The invention is also directed to use of the novel pyruvate compounds in reperfusion of tissue and organs both in vivo and in storage. Accordingly, the invention includes a method for the preservation of tissue deprived of oxygen
5 through events including, but not limited to, coronary infarction, stroke, mesenteric infarction, organ transplant (during preservation and intravenously after grafting of the organ) including amputated limbs. The compound is also believed well suited to treatment of acetaminophen poisoning
10 of the liver which depletes liver glutathione stores leading to acute hepatic necrosis.

This invention is also directed to the use of the novel pyruvate compounds to assist a patient's body in ethanol oxidation. In fact, the novel pyruvate compounds of this
15 invention are suited to use as nutritional supplements, preventing body fat deposition, lowering high blood cholesterol levels, and treatment for secondary diabetes effects. Furthermore, the novel pyruvates are believed superior for use in treating alcohol intoxication,
20 dermatological requirements and diabetic ketosis.

It is believed that the subject novel compounds provide stable, and physiological compounds with the beneficial result of delivering pyruvate and other NADH and ROS trapping moiety's.

25 Brief Description of the Drawings

The invention consists of the novel parts, construction and arrangements, combinations and improvements shown and described. The accompanying drawings, which are incorporated in and constitute a part of the specification illustrate one
30 embodiment of the invention and together with the description explain the principals of the invention.

Of the drawings:

FIGURE 1 shows swine infarct size after PNACE and DPAG treatment;

35 FIGURES 2, 3 and 4 show pyruvate ester blocking of UV inflammation;

FIGURE 5 shows pyruvate ester effect on UV pigmentation;

-8-

FIGURE 6 shows pyruvate ester relationship to inflammation;

FIGURE 7 shows DPAG effect on ketone bodies in ketoacidotic diabetic rats; and

5 FIGURE 8 shows DPAG effect on blood glucose in ketoacidotic diabetic rats.

Detailed Description of the Preferred Embodiments

For the purposes of this disclosure, the following abbreviations are used:

10 ADH, alcohol dehydrogenase; ALDH, aldehyde dehydrogenase; DCA, dichloroacetate; DHA, dihydroxyacetone; DHAP, dihydroxyacetone phosphate; DPAG, dipyruvyl-acetyl-glycerol; FAEE, fatty acid ethyl esters; GC, gas chromatography; GCMS, gas chromatography-mass spectrometry; LAD, left anterior
15 descending coronary artery; MS, mass spectrometry; NAC, N-acetylcysteine; NEFA, non-esterified fatty acids; PADA, pyruvyl-acetyl-dihydroxyacetone; PDAG, pyruvyl-diacetyl-glycerol; PDH, pyruvate dehydrogenase; PNACE, pyruvate N-acetylcysteine ethyl ester; ROS, reactive oxygen species.

20 PNACE

One preferred group of the inventive compounds is a thioester of pyruvate and a sulfur amino acid, for example cysteine or homocysteine. Preferably, any ionizable
25 removable radicals, such as ethyl and N-acetyl groups. The most preferred compound is formed of pyruvate and N-acetylcysteine ethyl ester.

Synthesis of PNACE

As understood in the art, pyruvate has proven to be a
30 relatively unstable compound with very limited mechanism for satisfactory delivery to subjects. However, the present inventive compound has proven to be readily manufacturable and very effective in the prevention of organ damage associated with reperfusion injury. The compound has been
35 prepared in pure form and in gram amounts. Its formula has

-9-

been confirmed by elemental analysis and gas chromatography-mass spectrometry. The compound is stable in slightly acidic solutions (pH 4-5). At pH 7.4, it is slowly hydrolyzed to pyruvate and N-acetylcysteine ethyl ester. The compound has
5 also been synthesized labeled with three deuterium ^2H atoms on the N-acetyl moiety. This deuterated compound is used as an internal standard for the assay of the compound by isotope dilution gas chromatography-mass spectrometry.

In a three-neck flask of 500 ml, freshly distilled
10 pyruvic acid (9.06 g., 0.102 mol) and N-hydroxy-succinimide (11.82 g., 0.102 mol) in dry tetrahydrofuran (THF) (180 ml) was stirred under nitrogen and was cooled in a ice bath. Dicyclohexylcarbodiimide (21.2 g., 0.102 mol) dissolved in dry THF (150 ml) was added slowly to the stirred cooled
15 mixture over approximately 1 hr. Then, the reaction mixture was stirred at room temperature for 2.5 hr, followed by slow addition of N-acetyl-L-cysteine ethyl ester (6.81 g., 0.033 mol) dissolved in 20 ml dry THF over approximately 1 hr. The reaction mixture was stirred overnight at room temperature
20 under a nitrogen atmosphere.

After evaporating the THF, the residue was suspended in ethyl acetate (750 ml) and was kept for 4-6 hr at 0°C. Dicyclohexyl urea (DCU) was then filtered and discarded; the ethyl acetate solution was washed three times with water
25 (3x100 ml). It was then dried over anhydrous sodium sulfate and concentrated under vacuum.

The product (17-18 g.) was further purified by using column chromatography. A column of 5 cm. diameter was filled with silica gel (180-200 g., 60 Angstrom flash chromatography
30 from Aldrich). The product was dissolved first in a minimum quantity of ethyl acetate:hexane (60:40) and was loaded on the column. The column was developed under gravity (rather than flash chromatography) with ethyl acetate:hexane (60:40). Fifty ml fractions were collected and monitored by TLC using
35 either iodine or UV light. The fractions containing the product were combined and solvents were removed under reduced pressure. The residue was dissolved in chloroform (300 ml), first washed with 5% HCl (2x30ml) and then saturated NaCl

-10-

(3x60 ml). The organic layer was dried over anhydrous sodium sulfate, filtered, and the solvent evaporated. The residue was dissolved in a minimum quantity of chloroform, and petroleum ether was added until the solution became turbid. 5 The suspension was kept overnight in the refrigerator and then filtered to get the pure crystallized product. The compound was dried under vacuum over P_2O_5 to a yield of 6.5 g. (75%, based on the N-acetyl-L-cysteine), m.p. 76-77°C.

Alternative Synthesis of PNACE

10 To a 250 ml three neck flask fitted with a thermometer, a magnetic stirrer, a 50-ml pressure-compensated addition funnel, and a Friedrich's condenser under nitrogen, was added 10g (52.3 mmol) of N-acetyl-L-cysteine ethyl ester, 8.0 ml of dry pyridine and 60 ml of dry benzene.

15 Pyruvoyl chloride (0.104 mole, 2 eq) was added dropwise over a period of 0.5 hr. while maintaining a temperature of 5°C to 10°C. Then, the reaction mixture was allowed to warm to 25°C and stirred for 2 hours at this temperature. The benzene solvent was then evaporated under vacuum. The crude 20 product was purified as above to yield 11.15 g of the desired compound (82%).

Synthesis of Deuterated PNACE

Pyruvate-N-[2H_3]acetyl-L-cysteine ethyl ester

Synthesized wherein the above procedure was followed 25 using N-[2H_3] acetyl-L-cysteine ethyl ester to form (d_3 -PNACE). The latter was prepared by reacting L-cysteine ethyl ester with [2H_6]acetic anhydride.

Set forth hereinbelow are certain analytical characteristics of the composition of the invention provided 30 to facilitate identification thereof, but not intended to limit the scope.

Characteristics of Compounds

I. Pyruvate-N-acetyl-L-cysteine ethyl ester : PNACE (unlabeled)

35 mp : 65°C

-11-

Rf (ethyl acetate/petroleum ether: 3/2) : 0.52

IR (Nicolet 300, CCl₄) (cm⁻¹) :

3435 (ν N-H)

3000 (ν C-H)

5 1747 (ν CO-O) ester

1731 (ν CO-S) thioester

1687 (ν CO-CO, CO-N) ketoester, amide

1497, 1378.3, 1210.1

10 NMR ¹H, 300MHz (Varian, CDCl₃, TMS) (ppm):1.33 (t, ³J=7.13, 3H, OCH₂CH₃)2.10 (s, 3H, COCH₃)2.50 (s, 3H, CH₃COCO)3.45 (dd, ³J=4.10 Hz, ³J=8.95 Hz, 2H, CH₂-S)4.23 (dd, ³J=7.13 Hz, 2H, CH₂CH₃)

15 4.83 (m, 1H, CH)

6.50 (sl, 1H, NH)

NMR ¹³C, 100.12 MHz (Bruker, CDCl₃, TMS)(ppm):

190.6; 188.08 keto, ketoester

168.6, 168.08 ester, amide

60.1 (OCH₂)49.5 (CH₂S)

28.4 (CHNH)

21.9 (CH₃COCO)20.8 (CH₃CO)12.1 (CH₃CH₂)

Mass spectrum, electron ionization (m/z):

20

190 (M - CH₃COCO, 33); 118(26);
102(56); 76(33), 60(90), 43 (CH₃CO⁺, 100)Mass spectrum, ammonia chemical
ionization (m/z):279 (M+18, 100); 262(M+1, 93); 209(49); 192(60)
175(18), 158(26)II. Pyruvate-N-[²H₃]acetyl-L-cysteine ethyl ester : d₃-PNACE
(deuterated)25 NMR ¹H, 300MHz (Varian, CDCl₃, TMS) (ppm) :1.34 (t, ³J=7.13, 3H, OCH₂CH₃)2.50 (s, 3H, CH₃COCO)3.42 (dd, ³J=4.10 Hz, ³J=8.95 Hz, 2H, CH₂-S)4.25 (dd, ³J=7.13 Hz, 2H, CH₂CH₃)

30 4.90 (m, 1H, CH)

6.50 (sl, 1H, NH)

NMR ¹³C, 100 MHz (Bruker, CDCl₃, TMS) (ppm):

190.5; 187.08 keto, ketoester

168.5, 168.10 ester, amide

60.1 (OCH₂)49.1 (CH₂S)

28.1 (CHNH)

20.8 (CH₃COCO)19.9 (CD₃CO)12.0 (CH₃CH₂)

Mass spectrum, electron ionization (m/z):


35

193 (M - CH₃COCO, 17); 121(4); 103(29);
77(12); 63(26); 43 (CH₃CO, 100)Mass spectrum, ammonia chemical
ionization (m/z):282(M+18, 42); 265(M+1, 47);
212(23), 195(37); 178(53); 161(100);
106(23); 89(15)40 Alternative, Suitable Pyruvate Ester Compounds

Also envisioned within the context of this invention are pyruvate ester compounds comprised of physiologically suitable sugars and polyols, including but not limited to ribose, galactose, glucose, fructose, sorbitol, inositol, arabitol, erythritol and other polyols. Within this group, ribose and glucose are particularly preferred.

-12-

Particularly esters of the formulae:



where R is H, pyruvyl or a short-chain acyl such as acetyl or propionyl and at least one R is pyruvyl.

5 DPAG and PADA

A further set of the inventive compounds are dipyruvyl-acetyl-glycerol (DPAG) and pyruvyl-acetyl-dihydroxyacetone (PADA). As with PNACE, these compounds are metabolizable substrates which counteract the effects of reperfusion injury. Glycerol is a physiological substrate which is well tolerated in large amounts and although DHA is not known to exist as such in body fluids, it is quickly phosphorylated by liver glycerol kinase to dihydroxyacetone phosphate (DHAP) which is a glycolytic intermediate. Similarly DPAG and PADA can be infused *in vivo* to deliver a therapeutic concentration of pyruvate without lactic acidosis and sodium overload. However, because DPAG and PADA can be administered in very high doses, they are also agents for accelerating ethanol oxidation in the liver, via transfer of reducing equivalents to peripheral tissues in the form of lactate.

Glycerol is a physiological substrate. It is released by adipose tissue lipolysis and is taken up by the liver, which is the major site of glycerol kinase activity (some glycerol kinase is also present in kidney). Glycerol kinase generates glycerol-phosphate which has 3 fates: glucose, glycerides/phospholipids, and lactate. DHA is converted to physiological dihydroxyacetone-phosphate (DHAP) by glycerol kinase. Then, DHAP has the same fates as glycerol. DHA is the oxidized counterpart of glycerol.

Because of the particular benefits of the thiol in PNACE, a dual strategy to prevent and/or treat reperfusion injury is considered advantageous. Moreover, to safely

-13-

deliver large amounts of pyruvate without sodium or nitrogen load, esters of pyruvate with either glycerol or dihydroxyacetone, i.e. DPAG or PADA, or the esters of sugars and other polyols described hereinbelow, can be used. PNACE is not a means to supply large amounts of pyruvate, since pharmacologically NAC concentrations of below 0.1 mM are often desirable, while effective pyruvate concentrations are 1- 2 mM. Thus, pyruvate-glycerol or pyruvate-DHA ester or the hereinbelow described sugar or polyol ester is infused in large amounts together with smaller amounts of PNACE.

Synthesis of DPAG

DPAG was prepared by esterification of 1-acetyl-glycerol (1-monoacetin) with pyruvyl chloride. To a 250 ml three-neck flask fitted with a thermometer, a mechanical stirrer, a 25 ml dropping funnel, and flushed with dry nitrogen, one adds 5.0 g of anhydrous monoacetin (dried for 2 days under vacuum), .0 ml of anhydrous pyridine, and 100 ml of anhydrous benzene. The flask is cooled below 10°C with an ice + salt slurry. Freshly distilled pyruvyl chloride (6.0 ml, 1 equivalent) is added dropwise over 15 min, while maintaining the temperature below 10°C. Then, the reaction mixture (showing a white precipitate of pyridinium chloride) is stirred for 1 hr at room temperature. The reaction mixture is filtered, to remove the pyridinium salt, and concentrated at 30°C on a rotavapor under high vacuum. The crude yellow product is dissolved in 50 ml of chloroform, washed once with 10 ml of HCl 1N, and stirred with 4g of Amberlyst-15 for 4 hr. The solvent is evaporated on a rotavapor under high vacuum at 30 °C maximum. The yield of DPAG (light yellow oil) is 9.6 g (94%).

The formula of DPAG was verified by (i) NMR ¹H and ¹³C, (ii) infrared spectra, (iii) enzymatic assay of the components of DPAG after hydrolysis, and (iv) HPLC before and after hydrolysis.

NMR ¹H (200 MHz Varian), solvent CDCl₃, reference TMS (δ in ppm):

-14-

5.30 (m, 1H, CH); 4.50-4.00 (m, 4H, CH₂O); 2.40 (s, 6H, CH₃COCO); 2.00 (s, 3H, CH₃CO)

NMR ¹H in agreement with the formula and the theoretical NMR spectra software ACD/LABS DEMO.

5 NMR ¹³C (200 MHz Varian), solvent CDCl₃, reference TMS (δ in ppm):

188.9, 188.7 (2C, carbonyls); 170.2 (1C, acetyl); 157.8, 157.9 (2C, pyruvyl); 68.9 (1C, CH); 61.5, 59.4 (2C, CH₂O); 24.6 (2C, CH₃); 18.4 (1C, CH₃) in agreement with the formula

10 and the theoretical NMR spectra software ACD/LABS DEMO.

IR (cm⁻¹, CCl₄) : 3537 (OH bonds from hydrated C=O), 2984; 1756, 1751, 1740, 1736, 1729 (C=O); 1383, 1231.

The NMR and IR spectra show that two molecules of water are fixed on carbonyl groups to form stable hydrated keto esters.

Incubation of DPAG with pig liver esterase liberates the components of the ester which were determined by enzymatic assays, thus confirming the formula of DPAG.

Synthesis of PDAG

20 PDAG was prepared in 81% yield by reacting diacetyl-glycerol with pyruvyl chloride, using the above procedure.

The formula of PDAG was verified by (i) NMR ¹H and ¹³C, (ii) infrared spectra, (iii) enzymatic assay of the components of DPAG after hydrolysis, and (iv) HPLC before and after hydrolysis.

NMR ¹H (200 MHz Varian); solvent: CDCl₃, reference TMS (δ in ppm):

5.28 (m, 1H, CH); 4.38-4.14 (m, 4H, CH₂O); 2.41 (s, 3H, CH₃COCO); 2.01 (s, 6H, CH₃CO). NMR ¹H in agreement with the

30 formula. NMR ¹³C (200 MHz Varian), solvent: CDCl₃, reference TMS (δ in ppm):

190.9 (1C, carbonyl); 170.4 (2C, acetyl); 159.8 (1C, pyruvyl); 71.5 (1C, CH acetyl); 68.6 (1C, CH pyruvyl); 61.9, 61.8 (2C, CH₂O); 24.7 (1C, CH₃ pyruvyl); 20.6 (2C, CH₃ acetyl)

35 in agreement with the formula. IR (cm⁻¹, CCl₄) : 3593 (OH bond, from hydrated C=O), 2973, 1762 (C=O bond), 1752 (C=O bond), 1744 (C=O bond), 1736 (C=O bond), 1374, 1242.

-15-

The NMR and IR spectra show that on a small fraction of the molecules, one molecule of water is fixed on a carbonyl group to form a stable hydrated keto ester.

Synthesis of PADA

5 PADA was prepared in 95% yield by esterification of dihydroxyacetone monoacetyl with pyruvyl chloride, as above.

TLC on silica (developed with chloroform/methanol/hexane 12/1/1 and revealed with iodine) showed one spot corresponding to PADA (Rf 0.45-0.50), and no
10 dihydroxyacetone, dihydroxyacetone monoacetate, or diacetate.

The formula of PADA was verified by (i) NMR ^1H and ^{13}C , (ii) infrared spectra, (iii) enzymatic assay of the components of PADA after hydrolysis, and (iv) HPLC before and after hydrolysis.

15 NMR ^1H (200 MHz Varian), solvent CDCl_3 , reference TMS (δ in ppm):

PADA (keto form) :4.94 (s, 2H, CH_2OCOCO); 4.74 (s, 2H, CH_2OCO); 2.49 (s, 3H, CH_3COCO); 2.08 (s, 3H, CH_3CO). NMR ^{13}C (200 MHz Varian), solvent CDCl_3 , reference TMS (δ in ppm):

20 PADA (keto form) :198.0 (1C, keto of DHA); 192.9 (1C, keto of pyruvyl); 170.1 (1C, acetyl); 159.1 (1C, pyruvyl); 67.3, 66.3 (2C, CH_2O); 26.7 (1C, CH_3 , pyruvyl); 20.3 (1C, CH_3 acetyl). Spectra in agreement with the formula. Enzymatic assay of pyruvate after hydrolysis was in agreement with the formula.

25 Reperfusion

As described in PCT/US97/04335, pyruvate is an effective drug for coronary reperfusion as treatment for acute myocardial infarction. The present compounds are equally suited for such an application.

30 Ethanol Metabolism

As fully described in PCT/US97/04335, ethanol can be exported from the liver, but there is no large-scale mechanism for exporting reducing equivalents from the liver. One obvious export mechanism would (i) trap reducing
35 equivalents in the conversion of pyruvate to lactate, and

-16-

(ii) export lactate to peripheral tissues. However, plasma pyruvate concentration is very low (0.05-0.1 mM). Pyruvate could be generated from glucose and amino acids, but these processes would further increase the liver's ATP burden. For these reasons several of the pyruvate compounds of the present invention are particularly suited to assist the body with ethanol oxidation.

Decrease in myocardial infarct size in swine by DPAG.

The animals were anesthetized, instrumented, and subjected to 60 minutes of total occlusion of the distal 2/3 of the left anterior descending coronary artery, followed by 120 min of reperfusion. The area of the infarct was stained with tetrazolium, and infarct size quantitated on 5 mm slices of tissue. Treatment with DPAG was started with the onset of reperfusion (8.0 mg/kg·min for 120 min). DPAG did not affect heart rate or ventricular pressure. Infarct size, expressed as a percent of the area at risk for infarction, was $40.0 \pm 3.6\%$ in the control group (n=9), which compares favorably with recently published values from another laboratory using the same swine model ($38.6 \pm 2.6\%$). With reference to Figure 1, the DPAG treated group (n=6) had an infarct size of $9.1 \pm 3.8\%$, which was statistically lower than the control group ($p < 0.0005$ by t-test). This demonstrated that DPAG, infused upon reperfusion, decreases markedly the size of myocardial infarct. A PNACE group (n = 4) had an infarct size of about $16 \pm 4\%$.

Acceleration of ethanol oxidation

Rat livers were infused with ethanol and the components of the esters of glycerol-and-DHA pyruvate to represent the conditions that will occur after ester hydrolysis. Livers from 24 h-fasted rats were perfused with non-recirculating buffer containing 4 mM glucose and 2 mM ethanol (20 times the K_m of ADH for ethanol, to insure zero order kinetics). After 10 min baseline, the influent perfusate was enriched with various equimolar concentrations of the components of the esters, ie DHA + Na-pyruvate, or glycerol + Na-pyruvate (up

-17-

to 2.2 mM). These conditions simulated infusion and hydrolysis of glycerol- or DHA-monopyruvate. The uptakes of ethanol, pyruvate, DHA and glycerol, as well as the productions of lactate and glucose were measured.

5 Addition of the components of the pyruvate esters increased ethanol uptake up to 5 fold (Figs 3 and 4). As expected, the uptake of ethanol was greater in the presence of DHA than in the presence of glycerol. This clearly shows that DHA contributes to the trapping of reducing equivalents
10 derived from ethanol oxidation. In perfusions with glycerol + pyruvate, correlation between pyruvate uptake and lactate output was linear with a slope of 0.7. Thus, 70% of the pyruvate taken up was converted to lactate. In perfusions with DHA + pyruvate, the correlation had also a slope of 0.7
15 up to a pyruvate uptake of 13 $\mu\text{mol}/\text{min}\cdot\text{g}$ dry wt (corresponding to influent DHA and pyruvate concentrations of 0.7 mM). At higher DHA and pyruvate concentrations, the slope increased to 1.45. However, at the highest DHA and pyruvate concentration used, the ratio (lactate
20 release)/(pyruvate uptake) was 0.96. The fraction of pyruvate uptake not accounted for was presumably converted to glucose and CO_2 . The uptake of glycerol and DHA increased with their concentration in the perfusate. As long as the (lactate production)/(pyruvate uptake) ratio was less than
25 1.0, there was no net conversion of glycerol or DHA to lactate. This occurred only at high DHA concentration. Thus, most of the glycerol and DHA were converted to glucose, glycerides, CO_2 or to a combination of these species. The relationship between ethanol uptake and lactate production.
30 Lactate yield was lower when pyruvate was infused with DHA rather than glycerol.

Before the infusion of the components of the pyruvate esters, the effluent [lactate]/[pyruvate] ratio could not be measured with precision, but must have been very high given
35 the presence of ethanol. As the concentrations of the ester components increased from 0.4 to 2.2 mM, the [lactate]/[pyruvate] ratio went down from about 12 to about 2. Thus, essentially all reducing equivalents generated from

-18-

ethanol were exported as lactate. The oxidized status of the liver NADH/NAD⁺ system may have allowed oxidation of part of the substrates, including acetate derived from ethanol.

In summary, these experiments confirmed that ethanol oxidation is stimulated by the components of DHA-pyruvate and glycerol-pyruvate. DHA is preferred as it acts not only as an esterifying group for pyruvate but also as a trap for reducing equivalents in its own right.

Therapeutic pyruvate concentration in vivo with DPAG

After preparing pure DPAG, experiments were performed to test whether it could be used to impose a therapeutic concentration of 1 mM pyruvate in arterial blood. Overnight-fasted rats, anesthetized with halothane, were infused in the jugular vein with DPAG at $90 \mu\text{mol} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$ for 90 min. This rate corresponds to about 120% of the rats' caloric requirement. Five blood samples (70 μl) were taken from the carotid artery between 60 and 90 min. The arterial concentrations of pyruvate, lactate, and glycerol were clamped at 1.0, 2.5, and 0.8 mM, respectively. Corresponding portal vein concentrations at 90 min were 0.6, 2.5, and 1.0 mM, respectively. Control rats show normal arterial concentrations of pyruvate (0.05 mM) and lactate (0.3 to 0.6 mM; normal values for lactate are up to 1.5 mM). In rats infused with DPAG, the arterial concentrations of pyruvate and lactate were clamped at 1.0, and 2.5 mM, respectively. Corresponding portal vein concentrations at 90 min were 0.6 and 2.5 mM, respectively. Arterial glucose remained at 5-6 mM. Final samples of aortic blood showed normal acid-base and electrolyte parameters. Thus, DPAG can be safely used to set up the 1 mM target concentration of pyruvate expected to be beneficial for the treatment of reperfusion injury. Similar data were obtained when PADA was infused to rats. This was achieved without sodium overload and/or acid-base perturbations. Second, the lack of major increases in glucose and lactate concentrations shows that administration of DPAG at 120% of the caloric requirement spares endogenous energy sources, probably including proteins. Third, during

-19-

peripheral administration of DPAG at $90 \mu\text{mol}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$, portal pyruvate concentration was about 2/3 that which yielded a 3 to 6-fold increase in ethanol uptake by perfused rat livers. A portal pyruvate concentration of 1 mM could be achieved (i) 5 by increasing the peripheral infusion of DPAG to $120 \mu\text{mol}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$, or (ii) by administering DPAG enterally to better target portal vein concentrations.

DPAG can thus be safely used to set up the 1 mM target concentration of arterial pyruvate expected to be beneficial 10 for the treatment of ethanol overdose and reperfusion injury. This was achieved without sodium overload and/or acid-base perturbations. Also, the lack of major increases in glucose and lactate concentrations shows that administration of DPAG at 120% of the caloric requirement spares endogenous energy 15 sources, probably including proteins.

The effect of DPAG on the rate of ethanol uptake by perfused rat livers was also tested. Livers were perfused with non-recirculating buffer containing 4 mM glucose, 2 mM ethanol and variable concentrations of DPAG (0 to 1.5 mM). 20 Uptake of ethanol by the liver increases 2.5 fold when DPAG concentration is raised from zero to 0.5 mM. Note that 0.5 mM DPAG corresponds to 1 mM pyruvate after hydrolysis. Thus, to accelerate ethanol oxidation *in vivo*, the rate of DPAG administration should be adjusted to achieve a 1 mM 25 concentration of free pyruvate in the portal vein. When DPAG was infused to live rats at $90 \mu\text{mol}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$, the portal vein concentration of pyruvate was 0.6 mM. A portal pyruvate concentration of 1 mM could be achieved *in vivo* (i) by increasing the peripheral infusion of DPAG to $120 \mu\text{mol}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$, or (ii) by administering DPAG enterally to 30 better target portal vein concentrations.

Accordingly, DPAG and PADA are effective in the treatment of alcoholic coma to prevent complications such as brain damage, hypothermia, respiratory depression, and 35 pulmonary infection and in the oral intake of the esters in conjunction with ingestions of alcoholic beverages, to accelerate ethanol oxidation and restore the capacity to drive a vehicle or operate machinery.

-20-

Pyruvate and the Krebs Cycle

As is well recognized, several oxidation steps in the Krebs cycle involve dicarboxylic and tricarboxylic acids. However, it is understood that a certain malfunction of this
5 cycle can occur via leakage of dicarboxylic and tricarboxylic acids (cataplerosis). It is believed the pyruvate delivered in accord with this invention can restore Krebs cycle effectiveness by "refilling" the pools of dicarboxylic and tricarboxylic acids (anaplerosis).

10 Dermatological uses of Pyruvate

Dermatologic applications of the above described pyruvate esters and thioesters have been identified. As described more fully herein below, the novel pyruvates can be used as active agents or as preservatives in topical and oral
15 dermatologic applications.

Particularly, the compounds are suitable for stabilization of all topical preparations or oral preparations to prevent oxidative damage to the ingredients in the formulation of, for instance, sunscreen chemicals and
20 their stabilizers, antioxidants and the associated preservatives, cosmetic antioxidants, bioactives, or other formulation purposes; oral agents in suspension in which the suspension needs or could be enhanced by a safe anti-oxidant.

Similarly, the compounds can be added to the composition
25 of tanning creams where the tanning effect of the dihydroxyacetone (DHA) moiety will be complemented by the antioxidant action of the pyruvate and thiol moieties of the pyruvate esters and thioesters.

The result is believed to be protection of the skin
30 against oxidative injury caused by a number of conditions including, but not limited to environmental stress from ultraviolet radiation (including prevention and treatment of sunburns) and routine suberythemogenic light exposure, and pollutants (including tobacco smoke), endogenous stress
35 resulting from diseases (cancer, infections, inflammatory

-21-

conditions, etc.), and diaper dermatitis-dermatological wound healing composition.

The compounds will also prevent, reduce and heal physical and chemical damage resulting from contact with harmful chemicals, radiation therapy damage including chemotherapy complication, laser exposure, laser surgery, electrochemical destruction, all skin peels, whether via chemical destruction (i.e., phenol, glycolic acids, alpha hydroxy acids, or other acids) or physical destruction (liquid nitrogen, laser light, other electromagnetic radiation or electro destructive, photodynamic therapies, etc.) can also be reduced via the present pyruvate compounds. In fact, the compound should be suitable to reduce general aging (chronologic aging), structural changes and wrinkling (photoaging, tobacco aging) of the skin.

In each instance, the pyruvate esters and thioesters of this invention are particularly advantageous because they are soluble in typical ointments and creams and less damaging to the skin than prior pyruvate compounds. Examples of suitable systems for delivery include ointments, creams, emulsions, lotions, mousses, gels, sprays, microencapsulated powders, solutions, dispersions, patches, bandages or any other form known to those skilled in the art.

Pyruvate esters, as an antioxidant, would be expected to attenuate inflammation, damage and aging of the skin through oxidative injury. Because pyruvate esters blocks UV-induced inflammation in human skin, one can reasonably predict that the esters will act on the inflammatory mechanisms that create inflammation in the skin after UV radiation injury, and which are shared by many other skin diseases. These include, but are not limited to, soluble mediator release (eicosanoids, histamine, mast cell products), reactive oxygen species mediated skin damage (both as a direct effect of photons reacting with skin cellular and structural elements and as a result of cellular ROS generated secondarily to the injury), DNA damage, cellular reactivity to UV-induced products (direct photoproducts, mediators, chromophores, DNA excision products, products of oxidation), cytokine release

-22-

and action, skin reconstruction, bone marrow derived cell activation, infiltration of leukocytes and their activation, vascular endothelial cell activation, adhesion molecule function and expression, fibroblast function, complement
5 activation, fibronectin modification, immune function alteration.

In addition, it is expected that the compounds will assist with all skin cell activations that occur in other skin conditions that result in generation of reactive oxygen
10 species or nitric oxide; most inflammatory reactions in the skin are associated with, and may be critically dependent upon, ROS or NO generation during the evolving pathogenic events in time and microanatomic location in the skin. In short, the pyruvate esters, as a safe and effective topical
15 antioxidant, would be useful as both a protecting ingredient and for promoting the health and resistance of skin against aging and injury. Similarly, it is believed that skin cell damage may be increased by the decrease of ROS by leukocytes which travel to the initially injured skin. The pyruvate
20 which are believed to block the leukocytes and/or their ROS generation can help to reduce this effect.

The following list of dermatological applications derives from the above understanding:

- Prevention and treatment of inflammation in the skin.
- 25 - Prevention of damage (including sunburn) from natural solar radiation or tanning beds or industrial/occupational exposure UV.
- Prevention of skin cancer, either via prevention of tumor promoting inflammation, or via prevention of oxidant
30 mediated mutations in DNA.
- Prevention of photoaging, such as loss of elasticity, immune function, development of solar lentigos (liver spots), actinic keratoses, dryness of the skin, wrinkling, sallow color.
- 35 - Prevention of skin injury and reduction of inflammation from chemicals, pesticides, wind or other environmental agents.
- Prevention of skin aging.

-23-

- Compensation for aging skin diminishment of ROS defense and the sequelae of such.

- Improve skin function in the face of repeated exposure to UV radiation and chemicals.

5 - Prevent skin aging processes attributable to ROS damage, including dryness, poor wound repair, poor immune function, easy bruisability, loss of elasticity, wrinkling, sallow color.

10 - Prevention and treatment of UV-induced immunosuppression and immune alteration of the skin

15 - Protection against injurious substances or environmental hazards which damage the skin such as carcinogens with direct or indirect oxidant effects, cold injury and frostbite, thermal and chemical burns, chemical irritants.

- Treatment of inflammatory skin diseases.

20 - Treatment of autoimmune diseases with activated leukocytes such as, but not limited to atopic dermatitis, xerotic eczema (dry skin), psoriasis, dermatitis herpetiformis, alopecia areata, granuloma annulare, sarcoid lupus erythematosus, lichen planus, scleroderma, graft vs. host disease.

25 - Treatment of other inflammatory diseases with activated leukocytes such as, but not limited to, contact dermatitis, drug reactions, erythema craquile, spongiotic dermatitis, lichen simplex chronicus, urticaria, toxic epidermal necrolysis, Stevens Johnson Syndrome, erythema multiforme.

30 - Treatment of other inflammatory skin conditions, such as, but not limited to, vasculitis, pyoderma gangrenosum, skin wounds, skin ulcers, keloids, scarring from wounding and wound repair.

35 - Prevention of exacerbation or relapse of inflammatory skin diseases that may be in partial or complete remission, such as, but not limited to, atopic dermatitis (eczema), psoriasis, xerotic eczemas, stasis dermatitis, psoriasis, lupus erythematosus.

-24-

- Reduction and prevention of vasculitis from all causes or no known underlying cause.

- Reduction and prevention of skin ulcers from all causes or no known underlying cause.

- 5 - Reduction of damage from melanomas and prevention of pigmentation response such as skin hyperpigmentation and early changing melanocytes.

It is believed that internal administration of compounds for prevention of photoinjury, thermal, or chemical
10 injury to areas not able to be protected by topical administration of material, for instance scalp, ears, lips, periocular skin, eyelids, conjunctivae, cornea, scleral conjunctivae, lens, vitreous, and retina and even the exposed skin diseases and conditions are certainly feasible. In any
15 event, it is believed that 1-30 ppm pyruvate may adequately achieve many of the above treatments. However, levels of about 0.01 to 1.0% and above are preferred.

Evaluations of Dermatological Applications

- 1) Pyruvate ester compounded in ethylene
20 glycol/aquaphor per separate notes at 0.01%, 0.03%, 0.1%, 0.3%, 1.0%, 3.0%

- 2) Stored at 4°C in capped syringes

- 3) Twice daily applications, beginning with 0.01% and observing the test site for irritation for 8-12 hours before
25 proceeding to the next concentration

- 4) 0.8 cc of each conc. was applied to the volar forearm at one of a set of premarked 2cm² areas corresponding to each conc.

- 5) After no irritation was observed for the 0-0.1%
30 preparations, the test sites were administered natural solar radiation (4 hr. post AM test material application)

- 6) Between 11:00 AM - 11:25 AM, the volar forearm was exposed to natural solar radiation, and 4 hr. later another 0.8cc of test materials were applied to the appropriate sites

- 35 6) 7 hours later the sites were graded by subjective assessment and Minolta chromameter.

-25-

Three chromameter measurements of three flashes each were taken from each site.

Results:

1) Irritation testing:

		Day	Redness	Surface	Sensation	Odor
5	Aquaphor	0	none	sl. xerosis	none	none
		1	none	smooth	none	none
		2	trace	smooth	none	none
10	0.01%	0	none	sl. xerosis	none	none
		1	none	smooth	none	none
		2	faint trace	smooth	none	none
	0.03%	0	none	sl. xerosis	none	none
		1	none	smooth	none	none
		2	none	smooth	none	none
15	0.10%	0	none	sl. xerosis	none	none
		1	none	smooth	none	none
		2	none	smooth	none	none
	0.30%	0	none	sl. xerosis	none	none
		1	none	smooth	none	none
		2				
20	1.00%	0	none	sl. xerosis	none	slight, pungent
		1	none	smooth	none	none
		2				

Redness: 7 hours post NSR

		Pyruvate Conc.	Chromameter Redness	Pyruvate Conc	Mean SEM	Redness: Subj.
25	Dry Adjacent Skin	0	9.74		9.74	trace
		0.00	10.17			trace
		0.00	9.42			
30	Aquaphor	0.00	10.50	0.00	10.03 0.32	
		0.01	8.72			faint - trace
		0.01	8.48			
35	mg% in Aquaphor	0.01	8.78	0.01	8.66 0.09	
		0.03	7.19			none
		0.03	6.81			
		0.03	6.20	0.03	6.73 0.29	
		0.10	6.51			none
		0.10	6.32			
40		0.10	6.41	0.10	6.41 0.05	

Conclusion:

Pyruvate ester in aquaphor is non-irritating and can block experimentally induced skin inflammation in human skin

-26-

		PIGMENTATION		48 hours post NSR	
		Pyruvate Ester Conc.	Chromameter pigmentation (1/L)	Pyruvate Conc.	pigmentation (1/L)
5		0	0.01523		
	Aquaphor	0.00			
		0.00		0.00	0.01523
	mg% in	0.01	0.01500		
10	Aquaphor	0.01		0.01	0.01500
	0.03	0.01476			
		0.03			
		0.03		0.03	0.01476
		0.10	0.01428		
15		0.10			
		0.10		0.10	0.1428
	dry, unexposed skin		0.01334		
20	Conclusion: Pyruvate ester prevented melanocytic response to UV as measured by tanning induction (reciprocal of lightness value by chromameter)				

REDNESS: 48 hours post NSR

		Pyruvate Ester Conc.	Chromameter Redness		Redness: Subj.
25	Dry Adjacent Skin	0	5.33		
	Aquaphor	0.00	9.40	5.33	trace
30		0.00		Pyruvate Conc	trace
		0.00		redness	
	mg% in	0.01	8.31	0.00	9.40
	Aquaphor	0.01			faint - trace
		0.01		0.01	8.31
35		0.03	6.28		none
		0.03		0.03	6.28
		0.10	6.94		none
		0.10		0.10	6.94

40 REDNESS: 48 hours post NSR

		Pyruvate Ester Conc.	Chromameter Redness	a*		
	Dry Adjacent Skin	0	5.33			5.33
45	Aquaphor	0.00	9.40	4.07	Pyruvate Conc.	redness
		0.00			0.00	9.40
	mg% in	0.01	8.31	2.98		
	Aquaphor	0.01				
50		0.01			0.01	8.31
		0.03	6.28	0.95		
		0.03				
		0.03			0.03	6.28
		0.10	6.94	1.61		
55		0.10				
		0.10			0.10	6.94

a* = Change in Redness from Baseline of Dry Adjacent Skin

-27-

These results are further represented in the Tables attached as Figures 2-6.

TREATMENT FOR DIABETIC KETOSIS

Under normal conditions, glucose and long-chain fatty acids (FA) are the major energy fuels for most tissues. Ketone bodies (KB), i.e., acetoacetate (AcAc) and R- β -hydroxybutyrate (R-BHB, also called D-BHB) are formed from the partial oxidation of FA in liver. These strong acids are totally ionized at physiological pH. KB are important fuels in fasting, strenuous exercise, postexercise recovery, etc (1,2). They accumulate in starvation and diabetic ketoacidosis (DKA).

With respect to this invention, it is believed in theory, but not a theory to which Applicants intend their invention to be limited that ketogenesis is fueled by FA released by adipose tissue lipolysis. The uptake of FA by the liver is directly proportional to their plasma concentration. In liver cytosol, FA are activated to acyl-CoAs which are either incorporated into lipids, or channeled to the mitochondria for oxidation. Transfer of acyl-CoAs to the mitochondria occurs via the carnitine palmitoyl transferases I and II (CPT I and II), with transient conversion to long-chain acyl-carnitines. The activity of CPT I, a major regulator of the entry of LC-acyl-CoAs into mitochondria, is inhibited by malonyl-CoA, an intermediate of fatty acid synthesis (see below). Once in liver mitochondria, acyl-CoAs undergo β -oxidation to acetyl-CoA. The three main fates of mitochondrial acetyl-CoA are (i) oxidation to CO₂ via citrate in the citric acid cycle (CAC), (ii) transfer to the cytosol via citrate and ATP-citrate lyase, and (iii) conversion to ketone bodies

Ketogenesis is regulated at three levels: supply of FA to the liver, β -oxidation of FA in liver, and energy status of the liver. First, the release of FA from adipose tissue is under hormonal control. It is stimulated by catecholamines and glucagon via hormone-sensitive lipase. It is inhibited by insulin. Second, in liver, the β -oxidation of FA is under

-28-

nutritional and hormonal control. It is activated by starvation, diabetes, and high-fat diet and inhibited by high-carbohydrate diet. In addition, FA oxidation is inhibited by malonyl-CoA, an intermediate of FA synthesis.

5 This provides a reciprocal control of FA synthesis and oxidation. Third, ketogenesis represents a spillover of carbon derived from FA β -oxidation which cannot be oxidized in the CAC. Since CAC activity is directly linked to ATP turnover, and since the flux through β -oxidation often

10 exceeds the CAC flux, excess β -oxidation carbon is exported as ketone bodies. This explains why hypermetabolic states (burn or crush injuries) that result in increased liver ATP turnover, are associated with low plasma ketone body concentrations in spite of the stimulation of lipolysis by

15 stress hormones.

Although ketogenesis can be greatly activated by dietary and hormonal manipulations, its absolute maximal rate is set by the O_2 uptake of the liver in a given metabolic situation. This rate can be calculated by assuming that all the O_2

20 uptake of the liver is used only to oxidize reducing equivalents formed during the conversion of FA to R-BHB.



For example, consider a 20 kg dog with a 500 g liver that takes up O_2 at $2 \mu\text{mol}\cdot\text{min}^{-1}\cdot(\text{g liver})^{-1}$ or $1 \text{ mmol}\cdot\text{min}^{-1}$, which

25 corresponds to a R-BHB production of $0.8 \text{ mmol}\cdot\text{min}^{-1}$, or $1.15 \text{ mol}\cdot\text{day}^{-1}$. This is clearly an unrealistic maximal rate since, to achieve it, no O_2 would be used for any other metabolic process in liver. However, this maximal rate is useful to discuss mechanisms of DKA.

30 R-BHB and AcAc represent a water-soluble form of FA which are transported from the liver to peripheral tissues. There, KB are converted to acetyl-CoA by mitochondrial 3-oxoacid CoA transferase (OAT) and AcAc-CoA thiolase:



35 $\text{AcAc-CoA} + \text{CoA} \leftrightarrow 2 \text{ acetyl-CoA (AcAc-CoA thiolase)}$

In peripheral tissues, acetyl-CoA derived from KB has two main fates, (i) oxidation to CO_2 via the CAC, and (ii) transfer to the cytosol via citrate and ATP-citrate lyase (in

-29-

lipogenic organs: adipose tissue, developing brain, lactating mammary gland).

In a given metabolic condition, the rate of KB uptake by peripheral tissues is roughly proportional to their plasma concentration over a wide dynamic range (1-4). When entering
5 a given tissue, KB compete with other fuels (FA and glucose) for the production of acetyl-CoA. This competition is influenced by the OAT activity of each tissue, the concentration of competing fuels, and the rate of acetyl-CoA
10 production from competing fuels

Consider the 20 kg dog. Its daily caloric requirement is $110 \cdot 20^{0.75} = 1040$ kcal/day. If all this caloric requirement were met by R-BHB oxidation (4.7 kcal/g), the dog would oxidize 221 g or 2.1 mol/day. This is about twice the
15 theoretical maximal rate of ketogenesis. There is evidence that the capacity of peripheral tissues to utilize KB is not far from the theoretical maximum. For example, dogs were infused with 1,3-butanediol diacetoacetate (4), a sodium-free precursor of KB (5), at 90% of their caloric requirement.
20 This corresponds to a rate of R-BHB infusion of 1.8 mol/day. The plasma concentration of total KB plateaued at 3 mM, reflecting metabolism of all the infused compound. Also, in pigs infused with 1,3-butanediol diacetoacetate at 30% of caloric requirement, the steady state total KB concentration
25 plateaued at only 0.5 mM. This is remarkable given that pigs are deficient in liver mitochondria HMG-CoA synthase, and have thus a very low capacity to make KB. Thus, in normal animals, peripheral tissues including the brain take up avidly KB when their plasma concentration is raised.

30 One of the main characteristics of decompensated insulin-dependent diabetes, in addition to hyperglycemia, hypokalemia, and dehydration, is ketoacidosis (up to 20 mM), resulting in life-threatening perturbations of acid/base status. DKA results from an imbalance between KB production
35 and utilization. Reviews from the literature emphasize the concept that DKA results mostly from hyperproduction of KB. Although a decrease in the capacity of peripheral tissues to utilize KB is mentioned. Strategies for decreasing ketosis

-30-

are solely aimed at (i) inhibiting adipose tissue lipolysis (insulin), and (ii) inhibiting the conversion of FA to KB in liver. To the best of our review of the literature, there are no strategies for accelerating KB utilization in peripheral
5 tissues. Based on the above, we think that underutilization of KB by peripheral tissues is a key component of the development of DKA.

To avoid the sodium load, we synthesized esters of pyruvate and glycerol or dihydroxyacetone (DHA). Glycerol and
10 DHA are the only esterifying compounds which can be administered in large amounts. Glycerol is a physiological substrate which is released by lipolysis mostly from adipose tissue. About 25% of glycerol production is taken up by the liver, 25% by the kidneys, and 50% by peripheral tissues.
15 Although DHA is not a physiological substrate, it is metabolized by glycerol kinase to DHA-phosphate, which is also a product of glycerol metabolism.

Prediabetic male BB rats were purchased from the University of Massachusetts Breeding Center. They were
20 treated with insulin (PZI 40 U/mL, Anpro Pharmaceutical), mean of 2.1 U for rats weighing between 250-300g) as soon as they became hyperglycemic. They were used after at least three weeks of clinical stability and steady weight growth. Under general anesthesia, permanent catheters were inserted
25 in a carotid artery and the contralateral external jugular vein. Then, the rats were deprived of insulin for two days. This resulted in hyperketonemia (9 -12 mM), hyperglycemia (22.6 ± 1.8 mM, n=18), and loss of weight (39g over 2 days). The rats were then treated with one of four protocols:
30 (i) saline infusion (0.267 ml/min. kg for 2h then 0.134 ml/min kg for 2h); (ii) saline infusion + 0.75 U of insulin intravenously (at 0 and 2 h); (iii) DPAG ($90 \mu\text{mol}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$, corresponding to 120% of caloric requirement of a normal rat of similar weight) in saline infusion; and (iv) DPAG in
35 saline infusion + 0.75 U of insulin intravenously (at 0 and 2 h).

Fig. 7 shows that, with DPAG treatment, total KB concentration decreased by 90% in 2 hr, much faster than with

-31-

insulin treatment (30%). When DPAG and insulin were administered together, their effect on ketosis was partly additive (Fig 7). Also, DPAG did not blunt the decrease in glucose concentration induced by insulin (Fig. 8). The data
5 show that massive doses of DPAG are well tolerated by compromised BB rats in diabetic ketoacidosis. Thus, we hypothesize that DPAG could become part of the treatment of DKA. DPAG may decrease ketosis by (i) inhibition of lipolysis in adipose tissue, (ii) inhibition of the conversion of FA to
10 KB in liver, and (iii) restoration of KB oxidation in peripheral tissues.

Pyruvate esters are not meant to replace the usual treatment of DKA by rehydration and insulin. They are meant to reinforce the usual treatment by accelerating the decrease
15 in KB concentration and the restoration of a normal blood pH (which is lowered by KB). With the usual treatment, the decrease in KB concentration to acceptable low values (less than 1 mM) takes from 8 to 15 hr. Based on our animal experiments, we think that ketosis could be resolved in 1 or
20 2 hr by infusing pyruvate esters at a rate ranging from 50 to 120% of the caloric requirement of the patient. The latter is calculated from the body weight by the formula

$$\text{Kcal/hr} = 4.6 \times (\text{body weight in kg})^{0.75}$$

Since the caloric density of DPAG and PADA is about 4
25 kcal/gram, the amount of pyruvate esters to be infused is about Grams pyruvate ester/hr = $0.9 \times (\text{body weight in kg})^{0.75}$

Thus it is apparent that there has been provided, in accordance with the invention, a pyruvate composition that fully satisfies the objects, aims, and advantages set forth
30 above. While the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications, and variations will be apparent to those skilled in the art in light of the foregoing description. Accordingly, it is intended to embrace all such
35 alternatives, modifications, and variations as fall within the spirit and broad scope of the appended claims.

-32-

We claim:

1. A method for treating or preventing dermatological illnesses or skin injury, reversing or slowing aging, or promoting skin health by aiding skin structure and function,
5 said method comprising orally, intravenously or topically administering a therapeutically effective amount of a pyruvate compound in the form of an ester of pyruvate and a sugar or a polyol, pyruvate thioester, or a dihydroxyacetone-pyruvate ester.
2. The method of claim 1 wherein said pyruvate compound is dipyruvyl-acetyl-glycerol.
3. The method of claim 1 wherein said pyruvate compound is pyruvyl-acetyl-dihydroxyacetone.
4. The method of claim 1 wherein said pyruvate compound is pyruvyl-diacetyl-glycerol.
5. The method of claim 1 wherein said pyruvate compound is a pyruvate thioester of cysteine or homocysteine.
6. The method of claim 1 wherein said pyruvate thioester is a N-acetyl derivative thereof.
7. The method of claim 1 wherein said prevention and treatment is directed to causations selected from the group consisting of radiation, cancer, photoaging, chemical, wind, cold, heat, autoimmune diseases, and inflammatory diseases.
8. The method of claim 1 wherein said pyruvate compound is glucose-pyruvate ester.
9. The method of claim 1 wherein said pyruvate compound is ribose-pyruvate ester.
10. A process for treating diabetic ketosis comprising administering an effective amount of an ester of pyruvate and

-33-

a sugar or a polyol, a pyruvate thioester, or a dihydroxyacetone-pyruvate ester.

11. The process of claim 10 wherein the pyruvate compound is dipyruvyl-acetyl-glycerol.

12. The process of claim 10 wherein said pyruvate compound is pyruvyl-acetyl-dihydroxyacetone.

13. The process of claim 10 wherein said pyruvate compound is pyruvyl-diacetyl-glycerol.


14. The process of claim 10 wherein said pyruvate compound is a pyruvate thioester of cysteine or homocysteine.

15. A method for administering pyruvate to a human which comprises treating said human orally, intravenously, or topically with an effective dosage or an ester of pyruvate and a sugar or a polyol.

16. The method of claim 15 wherein said sugar is further comprised of 4 to 7 carbon atoms.

17. The method of claim 15 wherein said sugar is selected from the group consisting of ribose, glucose and fructose.


18. The method of claim 17 wherein said pyruvate is of the formula:



where R is H, pyruvyl, or a short-chain acyl, and one to four R is pyruvyl.

-34-

19. The method of claim 17 wherein said pyruvate is of the formula:




where R is H, pyruvyl, or a short-chain acyl, and one to five R is pyruvyl.

20. The method of claim 13 wherein said polyol is selected from the group consisting of four to seven carbon polyols, aldoses or ketoses.

21. A topical formulation suitable for cosmetically or dermatologically administering to the skin, said formulation comprising a carrier and including a compound selected from the group consisting of pyruvate thioester, dihydroxyacetone-pyruvate, an ester of pyruvate and a sugar or a polyol and mixtures thereof.


22. The topical formulation of claim 21 wherein said compound is of the formula:




where one, two, or three R groups are pyruvyl and zero, one or two R groups may be a short-chain acyl selected from acetyl or propionyl.

-35-

23. The topical formulation of claim 21 wherein said compound is of the formula:




24. The topical formulation of claim 21 wherein said compound is of the formula:




where one or two R groups are pyruvyl and zero, one R group may be a short-chain acyl selected from acetyl or propionyl.

25. The topical formulation of claim 21 wherein said compound is of the formula:




-36-

26. The topical formulation of claim 21, wherein said compound is of the formula:




wherein R is selected from ethyl, methyl and alkyl groups.

27. The topical formulation of claim 21, wherein said compound is of the formula:



where R is H, pyruvyl, or a short-chain acyl, and where one to four R is pyruvyl.

28. The topical formulation of claim 21 wherein said compound is of the formula:




-37-

where R is H, pyruvyl, or a short-chain acyl, and where one to five R is pyruvyl.

29. The formulation of claim 21 comprised of between 1 and 30 ppm pyruvate.


30. The formulation of claim 21 comprised of at least 0.01% by weight pyruvate.

31. A compound for administering pyruvate to humans comprised of the formula:



where R is H, pyruvyl, or a short-chain acyl, and where one to five R is pyruvyl.


32. A compound for administering pyruvate to humans comprised of the formula:



-38-

where R is H, pyruvyl, or a short-chain acyl, and where one to five R is pyruvyl.

33. A compound for administering pyruvate to humans comprised of the formula:



where R is H, pyruvyl, or a short-chain acyl, and where one to four R is pyruvyl.

34. A method of stabilizing a cosmetic or dermatological composition comprising the step of including at least one esterified pyruvate.

1/8




FIG. 1




FIG. 2

3/8




FIG. 3

4/8

Pyruvate Ester Lowers UV Pigmentation

Change in Pigmentation from Baseline





FIG. 4

5/8

Pyruvate Ester Blocks UV Inflammation

**FIG. 5**

6/8




FIG. 6

7/8




FIG. 7

8/8





FIG. 8





INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/09729

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :A61K 7/48; 31/22 US CL :514/25, 858, 866, 893 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/25, 858, 866, 893 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WPIDS USPATFULL CAS ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,580,902 A (YU ET AL.) 03 December 1996, column 2, line 16 through column 7, line 34.	1-34
Y	5,602,183 A (MARTIN ET AL.) 11 February 1997, col. 5, line 26 through col. 40, line 59.	1-34
Y	4,970,143 A (GUIDOUX et al) 13 November 1990, col. 1, line 46 through col. 5, line 14.	21-34
Y	5,614,561 A (MARTIN) 25 March 1997, col. 10, line 32 through col. 17, line 40.	21-34
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* *A* *E* *L* *O* *P*	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* *X* *Y* *A* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
Date of the actual completion of the international search 03 AUGUST 1998		Date of mailing of the international search report 03 SEP 1998 
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer JAMES O. WILSON Telephone No. (703) 308-1235



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 413/14, 417/14, C07F 9/6558, C07D 413/12, 487/08, 451/02, 417/12, A61K 31/42, 31/44 // (C07D 487/08, 209:00, 209:00)</p>	A3	<p>(11) International Publication Number: WO 99/64417</p> <p>(43) International Publication Date: 16 December 1999 (16.12.99)</p>									
<p>(21) International Application Number: PCT/GB99/01753</p> <p>(22) International Filing Date: 3 June 1999 (03.06.99)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>9812021.5</td> <td>5 June 1998 (05.06.98)</td> <td>GB</td> </tr> <tr> <td>9820164.3</td> <td>17 September 1998 (17.09.98)</td> <td>GB</td> </tr> <tr> <td>9826066.4</td> <td>28 November 1998 (28.11.98)</td> <td>GB</td> </tr> </table> <p>(71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): GRAVESTOCK, Michael, Barry [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).</p> <p>(74) Agent: BRYANT, Tracey; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).</p>		9812021.5	5 June 1998 (05.06.98)	GB	9820164.3	17 September 1998 (17.09.98)	GB	9826066.4	28 November 1998 (28.11.98)	GB	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p> <p>(88) Date of publication of the international search report: 3 February 2000 (03.02.00)</p>
9812021.5	5 June 1998 (05.06.98)	GB									
9820164.3	17 September 1998 (17.09.98)	GB									
9826066.4	28 November 1998 (28.11.98)	GB									
<p>(54) Title: OXAZOLIDINONE DERIVATIVES, PROCESS FOR THEIR PREPARATION AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>(I)</p> </div> <div style="text-align: center;">  <p>(Q1)</p> </div> </div> <div style="display: flex; justify-content: space-around; align-items: flex-start; margin-top: 20px;"> <div style="text-align: center;">  <p>(Q2)</p> </div> <div style="text-align: center;">  <p>(TC5)</p> </div> </div> <p>(57) Abstract</p> <p>Compounds of formula (I), or a pharmaceutically-acceptable salt, or an in-vivo-hydrolysable ester thereof, wherein, for example, X is -O- or -S-; HET is an optionally substituted C-linked 5-membered heteroaryl ring containing 2 to 4 heteroatoms independently selected from N, O and S; Q is selected from, for example, formulae (Q1) and (Q2); R² and R³ are independently hydrogen or fluoro; T is selected from a range of groups, for example, an N-linked (fully unsaturated) 5-membered heteroaryl ring system or a group of formula (TC5) wherein R_c is, for example, R¹³CO-, R¹³SO₂- or R¹³CS-; wherein R¹³ is, for example, optionally substituted (1-10C)alkyl or R¹⁴C(O)O(1-6C)alkyl wherein R¹⁴ is optionally substituted (1-10C)alkyl; are useful as antibacterial agents; and processes for their manufacture and pharmaceutical compositions containing them are described.</p>											

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 99/01753

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D413/14 C07D417/14 C07F9/6558 C07D413/12 C07D487/08
C07D451/02 C07D417/12 A61K31/42 A61K31/44
//(C07D487/08,209:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 93 23384 A (THE UPJOHN CO.) 25 November 1993 (1993-11-25) cited in the application page 7, line 3 - line 21; claim 1 ---	1-6, 11-13
A	WO 97 09328 A (PHARMACIA & UPJOHN CO.) 13 March 1997 (1997-03-13) cited in the application page 16, line 15 - line 31; claim 1 --- -/--	1-6, 11-13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

30 November 1999

Date of mailing of the international search report

07/12/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 99/01753

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CHUNG-HO PARK ET AL: "ANTIBACTERIALS. SYNTHESIS AND STRUCTURE-ACTIVITY STUDIES OF 3-ARYL-2-OXOOXAZOLIDINES. 4 MULTIPLY-SUBSTITUTED ARYL DERIVATIVES" JOURNAL OF MEDICINAL CHEMISTRY, US, AMERICAN CHEMICAL SOCIETY. WASHINGTON, vol. 35, no. 6, 1992, page 1156-1165 XP000567006 ISSN: 0022-2623 cited in the application tables I, II</p> <p style="text-align: center;">---</p>	1-6, 11, 12
A	<p>W A GREGORY ET AL: "Antibacterials. Synthesis and Structure-Activity Studies of 3-Aryl-2-oxooxazolidines. 2. The A Group" JOURNAL OF MEDICINAL CHEMISTRY, US, AMERICAN CHEMICAL SOCIETY. WASHINGTON, vol. 33, 1990, page 2569-2578 XP002082548 ISSN: 0022-2623 cited in the application tables I-VII</p> <p style="text-align: center;">---</p>	1-6, 11, 12
A	<p>EP 0 645 376 A (MERCK PATENT GMBH) 29 March 1995 (1995-03-29) claim 1</p> <p style="text-align: center;">---</p>	1-6, 11
A	<p>EP 0 710 657 A (MERCK PATENT GMBH) 8 May 1996 (1996-05-08) claims 1, 5-7</p> <p style="text-align: center;">---</p>	1-6, 11
A	<p>WO 93 22298 A (TAIHO PHARMACEUTICAL CO LTD.) 11 November 1993 (1993-11-11) abstract</p> <p style="text-align: center;">---</p>	1-6, 11
A	<p>WO 94 22857 A (TAIHO PHARMACEUTICAL CO LTD.) 13 October 1994 (1994-10-13) abstract</p> <p style="text-align: center;">---</p>	1-6, 11
A	<p>WO 97 06791 A (SCRIPPS RESEARCH INST.) 27 February 1997 (1997-02-27) page 84, compound 15; page 87, compound 16; page 91, compound 17; page 94, compound 18</p> <p style="text-align: center;">-----</p>	1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 99/ 01753

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 10
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 10 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition (Rule 39.1 (IV) PCT-Method for treatment of the human or animal body by therapy).
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-5,9,11-13

Present claims 1-5, 9, 11-13 relate to an extremely large number of possible compounds and their preparation processes. The use of the term "optionally substituted" several times, without any further definition, in the claims makes the claims still more broad. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds of claims 6 to 8, the examples disclosed in the description, and their processes of preparation.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 99/01753

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9323384	A	25-11-1993	AU 668733 B	16-05-1996
			AU 4287793 A	13-12-1993
			CA 2133079 A	25-11-1993
			CN 1079964 A, B	29-12-1993
			CZ 9402505 A	16-08-1995
			EP 0640077 A	01-03-1995
			FI 945246 A	08-11-1994
			HU 72296 A	29-04-1996
			HU 9500659 A	28-11-1995
			IL 105555 A	15-07-1998
			JP 7506829 T	27-07-1995
			MX 9302665 A	01-11-1993
			NO 944237 A	04-01-1995
			PL 174909 B	30-10-1998
			PL 174850 B	30-09-1998
			SK 133794 A	07-06-1995
			US 5547950 A	20-08-1996
			US 5700799 A	23-12-1997
			ZA 9302855 A	24-10-1994
WO 9709328	A	13-03-1997	AU 6718196 A	27-03-1997
			BR 9610474 A	02-03-1999
			CA 2228647 A	13-03-1997
			CN 1197457 A	28-10-1998
			CZ 9800493 A	12-08-1998
			EP 0856002 A	05-08-1998
			FI 980452 A	27-02-1998
			NO 980855 A	30-04-1998
			PL 325152 A	06-07-1998
			SK 19598 A	04-11-1998
EP 645376	A	29-03-1995	DE 4332384 A	30-03-1995
			AT 178599 T	15-04-1999
			AU 682050 B	18-09-1997
			AU 7305094 A	06-04-1995
			CA 2132579 A	24-03-1995
			CN 1106806 A	16-08-1995
			CZ 9402247 A	12-07-1995
			DE 59408068 D	12-05-1999
			ES 2132295 T	16-08-1999
			HU 71233 A	28-11-1995
			JP 7179441 A	18-07-1995
			NO 943523 A	24-03-1995
			PL 305144 A	03-04-1995
			SK 112194 A	10-05-1995
			US 5561148 A	01-10-1996
			US 5723480 A	03-03-1998
			ZA 9407405 A	15-05-1995
EP 710657	A	08-05-1996	DE 19509093 A	09-05-1996
			AT 170179 T	15-09-1998
			AU 698987 B	12-11-1998
			AU 3452395 A	09-05-1996
			BR 9505039 A	21-10-1997
			CA 2161857 A	03-05-1996
			CN 1130626 A	11-09-1996
			CZ 9502858 A	15-05-1996
			DE 59503324 D	01-10-1998

INTERNATIONAL SEARCH REPORT

information on patent family members

Inter nal Application No

PCT/GB 99/01753

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 710657 A		ES 2123889 T	16-01-1999
		FI 955223 A	03-05-1996
		HU 74093 A	28-11-1996
		JP 8225550 A	03-09-1996
		NO 954366 A	03-05-1996
		PL 311194 A	13-05-1996
		SK 135895 A	05-06-1996
		TR 960383 A	21-06-1996
		ZA 9509212 A	24-07-1996
WO 9322298 A	11-11-1993	AU 653896 B	13-10-1994
		AU 4271093 A	29-11-1993
		CA 2112129 A,C	11-11-1993
		EP 0605729 A	13-07-1994
		FI 935920 A	29-12-1993
		HU 70423 A	30-10-1995
		HU 9500726 A	28-11-1995
		KR 146931 B	17-08-1998
		NO 934829 A	11-02-1994
		US 5480899 A	02-01-1996
WO 9422857 A	13-10-1994	AU 668818 B	16-05-1996
		AU 6437394 A	24-10-1994
		CA 2135474 A,C	13-10-1994
		EP 0645387 A	29-03-1995
		FI 945731 A	05-12-1994
		HU 72308 A	29-04-1996
		US 5521202 A	28-05-1996
WO 9706791 A	27-02-1997	AU 6846696 A	12-03-1997
		CN 1198667 A	11-11-1998
		CZ 9800409 A	16-09-1998
		EP 0844874 A	03-06-1998
		HU 9802675 A	29-03-1999
		JP 11511171 T	28-09-1999
		NO 980622 A	07-04-1998
		SK 18898 A	08-07-1998